

Quantum Equivalence of NC and YM Gauge Theories in 2 D and Matrix Theory

Badis Ydri^{*†}

February 2, 2008

Abstract

We construct noncommutative $U(1)$ gauge theory on the fuzzy sphere S_N^2 as a unitary $2N \times 2N$ matrix model. In the quantum theory the model is equivalent to a nonabelian $U(N)$ Yang-Mills theory on a 2 dimensional lattice with 2 plaquettes. This equivalence holds in the "fuzzy sphere" phase where we observe a 3rd order phase transition between weak-coupling and strong-coupling phases of the gauge theory. In the "matrix" phase we have a $U(N)$ gauge theory on a single point.

1 Introduction

A nonperturbative regularization of noncommutative gauge theory in two dimensions is obtained by putting the theory on a fuzzy sphere S_N^2 [1]. The lattice-like spacing parameter (the inverse UV cut-off) is proportional to $1/N$ where N is the size of the matrix algebra. In this regulator the radius of the sphere provides an IR cut-off for the theory. The limit $N \rightarrow \infty$ is the continuum limit [2].

The differential calculus on the fuzzy sphere is 3-dimensional and as a consequence a spin 1 vector field $\vec{A}$ is intrinsically 3-dimensional. Each component A_a , $a = 1, 2, 3$, is an element of Mat_N with $N = n(L + 1)$. Thus $U(n)$ symmetry will be implemented by $U(N)$ unitary transformations. On the fuzzy sphere S_{L+1}^2 it is not possible to split the vector field $\vec{A}$ in a gauge-covariant fashion into a tangent two-dimensional gauge field and a normal scalar fluctuation. We can only write a gauge-covariant expression for the normal field Φ as $\Phi = \frac{1}{2}(x_a A_a + A_a x_a + \frac{A_a^2}{\sqrt{c_2}})$ where x_a are the coordinates on fuzzy S_{L+1}^2 defined by $x_a = L_a / \sqrt{c_2}$ with $c_2 = \frac{L}{2}(\frac{L}{2} + 1)$. L_a are the generators of $SU(2)$ in the irreducible representation $\frac{L}{2}$. In the continuum limit $L \rightarrow \infty$ the normal scalar field reduces to $\Phi = n_a A_a$.

^{*}Email: ydri@physik.hu-berlin.de.

[†]This work is supported by a Marie Curie Fellowship from The Commission of the European Communities (The Research Directorate-General) under contract number MIF1-CT-2006-021797. The Humboldt-Universitat Zu Berlin preprint number is HU-EP-06/48.

The $U(n)$ action on the fuzzy sphere S_{L+1}^2 reads (with the identification $X_a = \alpha(L_a \otimes \mathbf{1}_n + A_a)$ and with the normalization $Tr_N \mathbf{1}_N = N$)

$$S = N \left[-\frac{1}{4} Tr_N [X_a, X_b]^2 + \frac{2i\alpha}{3} \epsilon_{abc} Tr X_a X_b X_c \right] + \frac{Nm^2}{2c_2} Tr_N (X_a^2 - \alpha^2 c_2)^2. \quad (1)$$

In the continuum limit this action becomes (with the gauge coupling constant defined by $g^2 = \frac{1}{N^2 \alpha^4}$)

$$S = \frac{1}{4g^2} \int \frac{d\Omega}{4\pi} tr_n F_{ab}^2 - \frac{1}{4g^2} \epsilon_{abc} \int \frac{d\Omega}{4\pi} tr_n [F_{ab} A_c - \frac{i}{3} [A_a, A_b] A_c] + \frac{2m^2}{g^2} \int \frac{d\Omega}{4\pi} tr_n \Phi^2. \quad (2)$$

If we take the limit $m \rightarrow \infty$ (in other words we set $\Phi = 0$) then the action will reduce to a 2-dimensional pure $U(n)$ gauge theory

$$S = \frac{1}{4g^2} \int \frac{d\Omega}{4\pi} tr_n F_{ab}^2 = \frac{1}{4g^2} \int \frac{d\Omega}{4\pi} tr_n \left(i\mathcal{L}_a A_b - i\mathcal{L}_b A_a + \epsilon_{abc} A_c + i[A_a, A_b] \right)^2. \quad (3)$$

In this last equation A_a should be understood as a 2-dimensional gauge field (in other words satisfying the constraint $n_a A_a = 0$) and $\mathcal{L}_a = -i\epsilon_{abc} n_b \partial_c$.

The model (1) with $m = 0$ was obtained in string theory limit in [3]. It was shown that it corresponds to a gauge theory on the fuzzy sphere in [4]. It was studied numerically in [8]. The model with $m = 0$ and $m \rightarrow \infty$ and for $U(1)$ groups was studied in one-loop perturbation theory in [7].

The basic prediction coming from the Monte Carlo study [5] is that noncommutative $U(1)$ gauge theory in 2 dimensions on the fuzzy sphere (given by the above action) behaves in the large N limit like a commutative $U(N)$ in two dimensions on a lattice [6]. This is true at least in the so-called "fuzzy sphere" phase of the model for large values of the gauge coupling constant. Indeed we observe in the simulation that this phase splits into two distinct regions corresponding to the weak and strong coupling phases of the gauge field which are separated by a third order phase transition. This transition seems to be consistent with that of a one-plaquette model [10]. It seems that deep inside the "fuzzy sphere" phase the model can still be understood as a $U(1)$ gauge theory on the sphere whereas in the "matrix" phase it is a $U(N)$ gauge theory on a single point.

The aim of this article is to construct models of noncommutative gauge theory on the fuzzy sphere which describe this third order phase transition although in general these models will coincide with (1) only in the continuum limit. An alternative approach to gauge theory on the fuzzy sphere is given in [9].

2 The Model

The main idea is to reparametrize the gauge field on S_N^2 in terms of a single matrix W which contains all the tangent degrees of freedom. This W we call the "fuzzy link variable" .

Thus we will have the coordinate transformation $(A_1, A_2, A_3) \longrightarrow (W, \Phi)$. Let us introduce the $2N \times 2N$ idempotent

$$\gamma = \frac{1}{N}(1 + 2\sigma_a L_a), \quad \gamma^2 = 1 \quad (4)$$

where σ_a are the usual Pauli matrices. It has eigenvalues $+1$ and -1 with multiplicities $N+1$ and $N-1$ respectively. We introduce the covariant derivative $D_a = L_a + A_a$ through a gauged idempotent γ_D as follows

$$\gamma_D = \hat{\gamma} \frac{1}{\sqrt{\hat{\gamma}^2}}, \quad \hat{\gamma} = \frac{1}{N}(1 + 2\sigma_a D_a), \quad \hat{\gamma}^2 = 1 + \frac{8\sqrt{c_2}}{N^2}\Phi + \frac{2}{N^2}\epsilon_{abc}\sigma_c F_{ab}. \quad (5)$$

In above $F_{ab} = i[D_a, D_b] + \epsilon_{abc}D_c = i[L_a, A_b] - i[L_b, A_a] + \epsilon_{abc}A_c + i[A_a, A_b]$. Clearly γ_D has the same spectrum as γ . Thus there exists a $U(2N)$ unitary transformation U such that $\gamma_D = U\gamma U^+$. The crucial observation is that if we let $T \in U(N+1) \times U(N-1)$ then $UT\gamma T^+ U^+ = U\gamma U^+$ since $T\gamma T^+ = \gamma$. In other words U is an element of the d_N -Grassmannian manifold $G_N = U(2N)/(U(N+1) \times U(N-1))$ and hence it (or equivalently γ_D) contains the correct number of degrees of freedom d_N which is found in a gauge theory on the fuzzy sphere without normal scalar field. Indeed $d_N = 4N^2 - (N+1)^2 - (N-1)^2 = 2N^2 - 2$. Thus γ_D can be identified with the equivalence class $[U] = \{UT, T \in U(N+1) \times U(N-1)\}$.

This fact can also be seen by writing down the measure in terms of U explicitly. A direct calculation gives $Tr_{2N}(d\gamma_D)^+(d\gamma_D) = 8 \sum_{i=1}^{N+1} \sum_{j=N+2}^{2N} (dU^+ U)_{ij} (U^+ dU)_{ji}$. It is obvious that only $d_N = 2N^2 - 2$ degrees of freedom of U are involved. In the large N limit we can see from (5) that $\gamma_D \longrightarrow \gamma$ and hence $U \longrightarrow \mathbf{1}_{2N}$. Thus in this limit we can write $U = 1 + i\frac{\Lambda}{N} + O(\frac{\Lambda^2}{N^2})$ and as a consequence

$$Tr_{2N}(d\gamma_D)^+(d\gamma_D) = \frac{8}{N^2} Tr_{N-1}(d\Lambda_{12})^+ d\Lambda_{12} + O(\frac{\Lambda^3}{N^3}). \quad (6)$$

In above Λ_{12} is the off-diagonal upper block of Λ and Λ_{12}^+ is the off-diagonal lower block of Λ . Λ_{12} is an $(N+1) \times (N-1)$ matrix while Λ_{12}^+ is an $(N-1) \times (N+1)$ matrix. We have then in the limit the measure

$$\int_{G_N} d\gamma_D \equiv \int_{G_N} d[U] \propto \int d\Lambda_{12} d\Lambda_{12}^+. \quad (7)$$

Integration over G_N means that we integrate over all idempotents γ_D which have $N+1$ eigenvalues equal $+1$ and $N-1$ eigenvalues equal -1 . A general idempotent with this property can be parametrized in terms of a unitary matrix U as $U\gamma U^+$. Thus (in the basis in which γ is diagonal with $\mathbf{1}_{N+1}$ in the first block and $-\mathbf{1}_{N-1}$ in the second block) we integrate only over the $d_N = 2N^2 - 2$ degrees of freedom of the unitary matrix U which correspond to γ_D . As we will see only these degrees of freedom will effectively appear in the action.

A general idempotent with the above property can also be parametrized in terms of 3 hermitian matrices D_a as in equation (5). Indeed we can check that we have (with γ_D given in terms of D 's by equation (5))

$$Tr_{2N}(d\gamma_D)^+ d\gamma_D = \frac{8}{N^2} Tr_N(dD_a)^+ dD_a - \frac{8}{N^2} Tr_N(d\Phi)^+ d\Phi + O(\frac{1}{N^3}). \quad (8)$$

Then in the large $N \rightarrow \infty$ limit it is obvious that we have the measure

$$\int dA_1 dA_2 dA_3 \equiv \int dD_1 dD_2 dD_3 \propto \int_{G_N} d\gamma_D \int d\Phi. \quad (9)$$

In the limit $m \rightarrow \infty$ we have $\Phi = 0$ and hence we obtain instead

$$\int dD_1 dD_2 dD_3 \delta(\Phi) \propto \int_{G_N} d\gamma_D. \quad (10)$$

Thus for consistency we must set $\Phi = 0$ in $\hat{\gamma}^2$. We can then compute (with the covariant coordinates x_a^D defined by $x_a^D = \frac{2}{N} D_a$ and $\epsilon\sigma F \equiv \epsilon_{abc}\sigma_c F_{ab}$) the following expansion

$$\gamma_D = \frac{1}{N} + \sigma x^D - \frac{1}{N^2}(\sigma x^D)(\epsilon\sigma F) - \frac{1}{N^3}(\epsilon\sigma F) + \frac{3}{2N^4}(\sigma x^D)(\epsilon\sigma F)^2 + \frac{3}{2N^5}(\epsilon\sigma F)^2 + O(\frac{1}{N^6}). \quad (11)$$

Let U and V be $2N \times 2N$ unitary matrices which are in the Grassmannian manifold G_N and let us consider the following path integral

$$Z[\gamma, \lambda] = \int_{G_N} d[U] \int_{G_N} d[V] \exp(-S_P), \quad S_P = -\frac{4N^2}{\lambda} + \frac{N}{\lambda} Tr_{2N} U \gamma U^+ V \gamma V^+ + \text{h.c.} \quad (12)$$

As we have explained integration over G_N means that we integrate only over the $2d_N = 2(2N^2 - 2)$ degrees of freedom of the unitary matrices U and V which will effectively appear in the action.

The matrix $U \gamma U^+$ can always be parametrized as $U \gamma U^+ = \gamma_D$ where γ_D is the idempotent given by (5). For small U near the identity given by $U = 1 + i\frac{\Lambda}{N} + \dots$ we can show that γ_D (and hence the action) will only depend on the off-diagonal blocks Λ_{12} and Λ_{12}^+ of Λ . We will only integrate over these components in the measure. Similarly the matrix $V \gamma V^+$ can be parametrized as $V \gamma V^+ = \gamma_B$ where γ_B is another idempotent given by equation (5) but in terms of a different covariant derivative $B_a = L_a + A'_a$ with a new gauge field A'_a and normal component Φ' . As before we write $V = 1 + i\frac{\Lambda'}{N} + \dots$ in the large N limit and again the action will only depend on the off-diagonal blocks Λ'_{12} and $(\Lambda'_{12})^+$ of Λ' . We will only integrate over these components in the measure.

The action S_P is invariant under all $U(2N)$ unitary transformations of the form $U \rightarrow gU$, $V \rightarrow gV$ which is only possible because of the doubling of gauge fields. The idempotents γ_D and γ_B will therefore transform as $\gamma_D \rightarrow g\gamma_D g^+$ and $\gamma_B \rightarrow g\gamma_B g^+$ respectively. Thus the measures $d\gamma_D$ and $d\gamma_B$ are also $U(2N)$ -symmetric and therefore the full path integral (12) is also $U(2N)$ -symmetric.

Next we compute the action S_P in the configuration $W = \gamma_D \gamma_B$. Let G_{ab} and x_a^B be the curvature and the covariant coordinates associated with the covariant derivative B_a . We obtain after a short calculation

$$\begin{aligned} S_P &= \frac{1}{N\lambda} Tr_N \left[-32c_2 + 2N^2(x_a^D + x_a^B)^2 + \frac{8i}{N^2}[D_a - B_a, D_b - B_b](F_{ab} + G_{ab}) - \frac{8}{N^2}(F_{ab} + G_{ab})^2 \right. \\ &+ \frac{6}{N^2}\{x_a^D, x_a^B\}(F_{cd}^2 + G_{cd}^2) - \frac{4}{N^2}x_a^D F_{cd} x_a^B G_{cd} - \frac{4}{N^2}x_a^B F_{cd} x_a^D G_{cd} \\ &+ \frac{2}{N^2}x_a^D (\epsilon F)_a x_b^B (\epsilon G)_b + \frac{2}{N^2}x_a^D (\epsilon F)_b x_b^B (\epsilon G)_a + \frac{2}{N^2}x_a^B (\epsilon F)_a x_b^D (\epsilon G)_b + \frac{2}{N^2}x_a^B (\epsilon F)_b x_b^D (\epsilon G)_a \\ &\left. - \frac{3}{N^2}([x_a^D, x_b^B] + [x_a^B, x_b^D])((\epsilon F)_a (\epsilon F)_b + (\epsilon G)_a (\epsilon G)_b) + O(\frac{1}{N^3}) \right]. \quad (13) \end{aligned}$$

These are the first few terms of the $U(1)$ action in this fuzzy plaquette model. The plaquette variable is the unitary $2N \times 2N$ matrix given by $W = \gamma_D \gamma_B$. This fuzzy plaquette $U(1)$ model is more complicated than the original fuzzy $U(1)$ model (1) although the two actions (1) and (13) have the same continuum limit as we will now show.

We show this result in two steps. First we compute the continuum limit of the above classical action then we integrate out one of the gauge fields. We will see in particular that the effect of the first non-trivial term $2N^2 \text{Tr}_N (x_a^D + x_a^B)^2$ is such that the path integration over B_a is dominated by the configuration $B_a = -D_a$. Indeed in the continuum large N limit in which we keep $N^2 \lambda$ fixed we can use in the classical action S_P the limits $x_a^D, x_a^B \rightarrow n_a$ where n_a are the global coordinates on the sphere and hence S_P reduces to

$$\begin{aligned} S_P &= -\frac{32c_2}{\lambda} + \frac{2N^2}{\lambda} \int \frac{d\Omega}{4\pi} (x_a^D + x_a^B)_*^2 + \frac{4}{N^2 \lambda} \int \frac{d\Omega}{4\pi} \left[F_{ab}^2 + G_{ab}^2 - 6F_{ab}G_{ab} + 2(\epsilon_{abc}n_c F_{ab})(\epsilon_{abc}n_c G_{ab}) \right] \\ &= -\frac{32c_2}{\lambda} + \frac{2N^2}{\lambda} \int \frac{d\Omega}{4\pi} (x_a^D + x_a^B)_*^2 + \frac{4}{N^2 \lambda} \int \frac{d\Omega}{4\pi} (F_{ab} - G_{ab})^2. \end{aligned} \quad (14)$$

In above we have used the fact that on the sphere $(\epsilon_{abc}n_c F_{ab})(\epsilon_{abc}n_c G_{ab}) = 2F_{ab}G_{ab}$ and $*$ stands for the star product on the fuzzy sphere which still appear in the second term. The path integral over the B_a is clearly seen to be dominated by the configuration $B_a = -D_a$ ¹. Therefore we obtain the action (modulo a constant term)

$$S_P = \frac{N}{\lambda} \text{Tr}_{2N} W + \text{h.c} = \frac{16}{N^2 \lambda} \int \frac{d\Omega}{4\pi} F_{ab}^2. \quad (15)$$

This action is essentially the $U(1)$ action obtained from (1) (see equation (3)) provided we make the following identification

$$\frac{16}{N^2 \lambda} \equiv \frac{1}{4g^2} \equiv \frac{\tilde{\alpha}^4}{4} \equiv \frac{\bar{\alpha}^4}{4N^2}. \quad (16)$$

Hence the fuzzy $U(1)$ action (1) with fixed coupling constant α corresponds in this particular limit to the fuzzy $U(1)$ plaquette action (13) in the weak regime $\lambda \rightarrow 0$ and agreement between the two is expected only for weak couplings (small values of λ or equivalently large values of $\bar{\alpha}$). This is what we observe numerically [5].

3 The solution

The path integral (12) is invariant under all $U(2N)$ unitary transformations of the form $U \rightarrow gU$, $V \rightarrow gV$ or equivalently $\gamma_D \rightarrow g\gamma_D g^+$, $\gamma_B \rightarrow g\gamma_B g^+$. This symmetry can be fixed as follows. We perform the following $U(2N)$ transformation $U \rightarrow U' = gU$ with $g = V^+$. For fixed V we obtain the measure $\int_{G_N} dU = \int_{G_N} dU'$ and the action S_P will only depend on U' .

¹This means in particular that the unitary matrix $W = \gamma_D \gamma_B$ approaches in the quantum theory $-\mathbf{1}_{2N}$ in contrast with the classical limit $W \rightarrow \mathbf{1}_{2N}$.

The integral over V can be done and one ends up with the path integral (by dropping also the primes)

$$\begin{aligned} Z[\gamma, \lambda] &= \int_{G_N} d[U] \exp \left(\frac{4N^2}{\lambda} - \frac{N}{\lambda} \text{Tr}_{2N}(\gamma U \gamma U^+ + \text{h.c.}) \right) \\ &\equiv \int_{G_N} d\gamma_D \exp \left(\frac{4N^2}{\lambda} - \frac{N}{\lambda} \text{Tr}_{2N}(\gamma \gamma_D + \text{h.c.}) \right). \end{aligned} \quad (17)$$

This is still invariant under $U(N+1) \times U(N-1)$ unitary transformations $U \rightarrow gU$. This is the gauge symmetry we want on the fuzzy sphere with $N \times N$ matrices. Let us recall that we want a $U(N)$ gauge symmetry on the fuzzy sphere but in this construction where we are using $2N \times 2N$ matrices it is only natural to obtain a gauge group which is twice as large. Remark also that we do not get precisely $U(N)$ but the groups $U(N+1)$ and $U(N-1)$. In the large N limit this becomes unimportant.

Apart from the restricted integration which is performed on the Grassmannian manifold G_N the path integral (17) looks very much like a $U(2N)$ one-plaquette model. Indeed the matrix $W = \gamma \gamma_D$ is a unitary $2N \times 2N$ matrix. Let us then write (17) in the equivalent form

$$\begin{aligned} Z[\gamma, \lambda] &= \int dW \int_{G_N} d\gamma_D \delta(W - \gamma \gamma_D) \exp \left(\frac{4N^2}{\lambda} - \frac{N}{\lambda} \text{Tr}_{2N}(W + \text{h.c.}) \right) \\ &= \int dW I(W) \exp \left(\frac{4N^2}{\lambda} - \frac{N}{\lambda} \text{Tr}_{2N}(W + \text{h.c.}) \right). \end{aligned} \quad (18)$$

We can therefore approximate the partition function of noncommutative $U(1)$ gauge theory on the fuzzy sphere given by (12) with the following path integral

$$\begin{aligned} Z[\gamma, \lambda, M^2] &= \int dW I(W, M^2) \exp \left(\frac{4N^2}{\lambda} - \frac{N}{\lambda} \text{Tr}_{2N}(W + \text{h.c.}) \right) \\ I(W, M^2) &= \int_{G_N} d\gamma_D \exp \left(-M^2 \text{Tr}_{2N}(W - \gamma \gamma_D)^+ (W - \gamma \gamma_D) \right). \end{aligned} \quad (19)$$

Instead of the delta function in (18) which implements the constraint $W - \gamma \gamma_D = 0$ we add a new term to the action with large positive coupling constant M^2 which implements this constraint only approximately. As it turns out the integral over G_N in (19) is much easier to compute (at least in large N) than the original corresponding path integral $I(W)$ in (18).

We remark that by dropping the requirement that M^2 must be large we get a model with an enlarged phase space. Indeed the space of parameters of the theory becomes the two dimensional quadrant $\lambda \geq 0, M^2 \geq 0$ instead of the original positive real line. An alternative way of thinking about (19) is as follows. The one-parameter family of models $Z_M \equiv Z[\gamma, \lambda, M^2]$ interpolate between the model Z_0 which corresponds to a true $U(2N)$ one-plaquette model and Z_∞ which is the path integral of the noncommutative $U(1)$ gauge theory on the fuzzy sphere given by (18) or equivalently (12). In other words holding N fixed and taking the limit $M^2 \rightarrow \infty$ reproduces the original path integral (18).

It is hoped that the models Z_M with intermediate large values of M^2 will capture some of the most important features of noncommutative $U(1)$ gauge theory on the fuzzy sphere (at

least in the "fuzzy sphere" phase) seen in the Monte Carlo study of the action (1). In particular we will consider in the following the double scaling limits $M^2, N \rightarrow \infty$ keeping M^2/N fixed. By construction the models Z_M coincide in these double scaling limits with the limit $N \rightarrow \infty$ of (18). So these models have the correct continuum limit and hence they are also path integrals of noncommutative $U(1)$ gauge theories on the fuzzy sphere. However quantum mechanically these models for finite fixed values of the ratio M^2/N are found to behave differently from (18) which corresponds to the ratio $M^2/N \rightarrow \infty$. As we will see the limit $M^2/N \rightarrow 0$ is also different.

The matrix $\gamma\gamma_D$ is a unitray matrix which in the continuum large N limit tends (by equation (5)) to the identity matrix $\mathbf{1}_{2N}$. Indeed in this limit both γ and γ_D approach the usual chirality operator $\gamma = n_a \sigma_a$ and hence $\gamma\gamma_D \rightarrow \mathbf{1}_{2N}$. Equivalently the unitary matrix U will approach in the continuum limit the identity $\mathbf{1}_{2N}$ as $U = 1 + i\frac{\Lambda}{N} + O(\frac{\Lambda^2}{N^2})$. In this limit it is also seen by equation (7) that the measure $\int_{G_N} d\gamma_D \equiv \int_{G_N} d[U]$ depends only on the off-diagonal blocks Λ_{12} and Λ_{21} with $\Lambda_{21}^+ = \Lambda_{12}$. Let W_1 and W_2 be the diagonal blocks of W in the basis where γ is diagonal. We denote by W_{12} and W_{21} the off-diagonal blocks. Thus in the large N limit the second line of (19) takes the form

$$\begin{aligned} I(W, M^2) &= \int_{G_N} d\gamma_D e^{-M^2 \text{Tr}_{2N} (2 - W^+ \gamma \gamma_D - W \gamma_D \gamma)} \\ &\propto \int d\Lambda_{12} d\Lambda_{21} e^{-M^2 \text{Tr}_{2N} (W-1)^+ (W-1) - \frac{2iM^2}{N} \text{Tr}_{N+1} (W_{21}^+ - W_{12}) \Lambda_{21} - \frac{2iM^2}{N} \text{Tr}_{N-1} (W_{12}^+ - W_{21}) \Lambda_{12}}. \end{aligned} \quad (20)$$

There are two cases to consider.

Case I : This corresponds to the special case $M^2 = 0$. By the requirement of continuity this point should be reached in the double scaling limits $M^2, N \rightarrow \infty$ keeping $\frac{M^2}{N}$ fixed such that $\frac{M^2}{N} \rightarrow 0$. The above integral (20) becomes

$$I(W, M^2) = e^{-M^2 \text{Tr}_{2N} (W-1)^+ (W-1)}. \quad (21)$$

Thus we end up with the partition function

$$Z[\gamma, \lambda, M^2] \propto \int dW \exp \left(\frac{4N^2}{\lambda_M} - \frac{N}{\lambda_M} \text{Tr}_{2N} (W + \text{h.c.}) \right), \quad \frac{1}{\lambda_M} = \frac{1}{\lambda} - \frac{M^2}{N}. \quad (22)$$

Case II : This is the generic case of the double scaling limits $M^2, N \rightarrow \infty$ keeping $\frac{M^2}{N}$ finite and fixed. Extrapolation of the results to the case $\frac{M^2}{N} \rightarrow \infty$ will by definition correspond to the model (18). However extrapolation of the results to $\frac{M^2}{N} \rightarrow 0$ will not reproduce case I as we will see. In these double scaling limits the integral (20) becomes

$$I(W, M^2) \propto e^{-M^2 \text{Tr}_{N+1} (2 - W_1 - W_1^+) - M^2 \text{Tr}_{N-1} (2 - W_2 - W_2^+)} \delta(W_{21}^+ - W_{12}) \delta(W_{12}^+ - W_{21}). \quad (23)$$

In above W_1 is an $(N+1) \times (N+1)$ matrix, W_2 is an $(N-1) \times (N-1)$ matrix and W_{12}, Λ_{12} are $(N+1) \times (N-1)$ matrices whereas W_{21}, Λ_{21} are $(N-1) \times (N+1)$ matrices.

Since in these limits M^2 is large proportional to N we also see that we have $W \rightarrow \mathbf{1}_{2N}$ or equivalently $W_1 \rightarrow \mathbf{1}_{N+1}$ and $W_2 \rightarrow \mathbf{1}_{N-1}$ and hence $W_{12} + W_{21}^+ \rightarrow 0$ which follows from the identity $W_1^+ W_{12} + W_{21}^+ W_2 = 0$. Together with the delta function $\delta(W_{21}^+ - W_{12})\delta(W_{12}^+ - W_{21})$ appearing in the last line of equation (23) we can conclude that the off-diagonal blocks W_{12} and W_{21} of W will approach zero in this double scaling limit. By neglecting also the edge effects in the large N limit we can take W_1 and W_2 to be $N \times N$ matrices. The partition function given by the first line of (19) becomes

$$Z[\gamma, \lambda, M^2] \propto [Z(\lambda_M)]^2, \quad Z(\lambda_M) = \int dW_1 \exp\left(\frac{2N^2}{\lambda_M} - \frac{N}{\lambda_M} \text{Tr}_N(W_1 + \text{h.c.})\right). \quad (24)$$

Therefore in these double scaling continuum limits in which $M^2, N \rightarrow \infty$ keeping M^2/N fixed we can set $I(W)$ in the first line of equation (19) equal to 1 and replace W with the matrix obtained by taking the diagonal parts W_1 and W_2 to be two arbitrary independent unitary $N \times N$ matrices while allowing the off-diagonal parts W_{12} and W_{21} to go to zero. In this approximation we can drop the restriction that these unitary matrices W_1 and W_2 must be close to $\mathbf{1}_N$ since we have replaced the coupling constant λ with λ_M .

The path integral of a 2-dimensional $U(N)$ gauge theory in the axial gauge $A_1 = 0$ on a lattice with volume V and lattice spacing a is given by $Z(\lambda_M)^{V/a^2}$. Therefore we can see that the partition function $Z(\gamma, \lambda, M^2)$ of noncommutative $U(1)$ gauge theory on the fuzzy sphere is proportional to the partition function of a 2-dimensional $U(N)$ gauge theory in the axial gauge $A_1 = 0$ on a lattice with two plaquettes. This is true at least in the weak coupling region of the phase space. Indeed since $\lambda_M \geq 0$ we must have

$$\lambda \leq \frac{N}{M^2}. \quad (25)$$

The matrix phase The fact that the one-plaquette model (24) can be defined for all values of the coupling constant λ_M in the range $[0, \infty[$ leads to the above restriction on the allowed values of the gauge coupling constant λ . This can be understood as follows. The partition function $Z(\gamma, \lambda, M^2)$ corresponds to noncommutative $U(1)$ gauge theory on the fuzzy sphere only for values of the coupling constant λ below the upper value $\frac{N}{M^2}$ where we can make sense of this model as a one-plaquette model. For $\lambda > \frac{N}{M^2}$ the model $Z(\gamma, \lambda, M^2)$ is presumably in the so-called "matrix" phase where there is no an underlying stable sphere as a spacetime and the gauge theory is defined on a single point.

This picture is consistent with our previous results [5, 7]. Indeed we found in the one-loop calculation as well as in numerical simulation that the model (1) undergoes a first order phase transition from the "fuzzy sphere" phase to a "matrix" phase where the fuzzy sphere vacuum collapses under quantum fluctuation. The fuzzy sphere phase is defined only for values of the coupling constant λ such that (m^2 is the mass parameter appearing in (1))

$$\lambda \leq \lambda_*^{S^2 \rightarrow \{0\}} = \frac{8}{N^2}(m^2 + \sqrt{2} - 1). \quad (26)$$

The one-plaquette 3rd order phase transition In this section we solve the models given by (22) and (24) in the large N limit. The only difference between the two models lies in the fact that (22) is a $U(2N)$ plaquette model while (24) is a $U(N)$ plaquette model and hence we will concentrate on the detail of (24) which describes noncommutative $U(1)$ gauge theory on the fuzzy sphere.

We can immediately diagonalize the matrix W_1 in (24) by writing $W_1 = TDT^+$ where T is some $U(N)$ matrix and D is diagonal with elements equal to the eigenvalues $\theta_i + \pi$ of W_1 ². In other words $D_{ij} = \delta_{ij}(\theta_i + \pi)$. The integration over T can be done trivially and one ends up with the path integral

$$Z(\lambda_M) = \int \prod_{i=1}^N d\theta_i e^{NS_N}, \quad S_N = \frac{2N}{\lambda_M} + \frac{2}{\lambda_M} \sum_{i=1}^N \cos \theta_i + \frac{1}{2N} \sum_{i \neq j} \ln \left(\sin \frac{\theta_i - \theta_j}{2} \right)^2. \quad (27)$$

The last term is due to the Vandermonde determinant. In the large N limit we can resort to the method of steepest descent to evaluate the path integral $Z(\lambda)$. The partition function will be dominated by the solution of the equation $\frac{dS_N}{d\theta_i} = 0$ which is a minimum of the action S_N . In the large N limit we introduce a density of eigenvalues $\rho(\theta)$ which is positive definite and normalized to one. Thus sums will be replaced by $\sum_i = N \int d\theta \rho(\theta)$. The saddle point solution must satisfy the equation of motion

$$\frac{2}{\lambda_M} \sin \theta = \int d\tau \rho(\tau) \cot \frac{\theta - \tau}{2}. \quad (28)$$

By using the expansion $\cot \frac{\theta - \tau}{2} = 2 \sum_{n=1}^{\infty} (\sin n\theta \cos n\tau - \cos n\theta \sin n\tau)$ we can solve this equation quite easily in the strong-coupling phase and one finds the solution [10]

$$\rho(\theta) = \frac{1}{2\pi} + \frac{1}{\pi\lambda_M} \cos \theta. \quad (29)$$

However it is obvious that this solution makes sense only where the density of eigenvalues is positive definite. This density of eigenvalues ρ is positive definite for all values of θ iff $\lambda_M > 2$. The critical value of the coupling constant λ_M is then seen to occur at $\lambda_{M*} = 2$. In terms of the coupling constant λ we obtain the critical value

$$\lambda_*^u = \frac{2}{1 + 2\frac{M^2}{N}}. \quad (30)$$

From (25) we then see that for the weak coupling phase we have $\lambda \leq \lambda_*^u \leq \frac{N}{M^2}$. In the strong coupling phase $\lambda_*^u \leq \lambda \leq \frac{N}{M^2}$.

In the weak regime we obtain the solution [10]

$$\rho(\theta) = \frac{2}{\pi\lambda_M} \cos \frac{\theta}{2} \sqrt{\frac{\lambda_M}{2} - \sin^2 \frac{\theta}{2}}, \quad \sin \frac{\theta_*}{2} = \sqrt{\frac{\lambda_M}{2}}. \quad (31)$$

²The addition of the angle π reflects the fact that the unitary matrix W approaches in the quantum theory $-\mathbf{1}_{2N}$ and not $\mathbf{1}_{2N}$ for very weak couplings. See the footnote on page 5.

Again the critical value of the coupling constant λ_M is seen to occur at $\lambda_{M*} = 2$ when the critical angle θ_* approaches π . At this point the eigenvalues θ of W_1 will fill the whole unit circle.

For very strong couplings the density of eigenvalues (29) becomes a uniform distribution while in the very weak coupling region (which corresponds to very small angles θ) the density of eigenvalues (31) reduces to Wigner semicircle law, viz

$$\rho(\theta) = \frac{1}{\pi\lambda_M} \sqrt{2\lambda_M - \theta^2}. \quad (32)$$

We can use this last equation to compute the free energy and specific heat for small values of the coupling constant λ with excellent agreement with the simulation results of (1). The free energy is given by

$$\begin{aligned} \frac{F_w}{N^2} = 2 \times \frac{S_N}{N} &= \frac{8}{\lambda_M} - \frac{2}{\lambda_M} \int_{-\theta_*}^{\theta_*} d\theta \rho(\theta) \theta^2 + \int_{-\theta_*}^{\theta_*} d\theta \rho(\theta) \int_{-\theta_*}^{\theta_*} d\alpha \rho(\alpha) \ln \left(\frac{\theta - \alpha}{2} \right)^2 \\ &= \frac{8}{\lambda_M} - 1 + \ln \frac{\lambda_M}{2} + c_1. \end{aligned} \quad (33)$$

The factor of 2 which multiplies $\frac{S_N}{N}$ comes from the fact that we have two identical independent one-plaquette models contributing to F in accordance with (24). In order to compute the specific heat we implement the scaling transformations $F \rightarrow \frac{F}{T}$ and $\lambda_M \rightarrow T\lambda_M$. The specific heat is then defined by

$$C_v = \left(T^3 \frac{\partial^2 F}{\partial T^2} \right)_{T=1}. \quad (34)$$

A straightforward calculation yields the very simple result

$$\frac{C_v}{N^2} = 1. \quad (35)$$

This is precisely what we see numerically in the fuzzy sphere phase. The value 1 emerges from the fact that we have two plaquettes. As it turns out this result is valid throughout the weak-coupling phase, i.e for all $\lambda_M \leq 2$.

In the regime of strong couplings the free energy and specific heat are computed using the distribution of eigenvalues (29). We find

$$\begin{aligned} \frac{F_s}{N^2} = 2 \times \frac{S_N}{N} &= \frac{4}{\lambda_M} + \frac{4}{\lambda_M} \int_{-\pi}^{\pi} d\theta \rho(\theta) \cos \theta + \int_{-\pi}^{\pi} d\theta \rho(\theta) \int_{-\pi}^{\pi} d\alpha \rho(\alpha) \ln \left(\sin \frac{\theta - \alpha}{2} \right)^2 \\ &= \frac{4}{\lambda_M} + \frac{4}{\lambda_M^2} + \frac{4}{\pi^2 \lambda_M^2} \int_0^{\pi} dx (\pi - x) \cos 2x \ln(\sin x)^2 + c_2 \\ &= \frac{4}{\lambda_M} + \frac{2}{\lambda_M^2} + c_2. \end{aligned} \quad (36)$$

The specific heat in this phase is therefore given by

$$\frac{C_v}{N^2} = \left(\frac{2}{\lambda_M} \right)^2. \quad (37)$$

The phase diagram From equation (30) we can immediately see that for every value of the parameter M^2/N the model (24) undergoes a third order phase transition from strong-coupling phase to weak-coupling phase consistent with the ordinary one-plaquette third order phase transition observed in 2 dimensional gauge theory on the lattice [10]. However in this case this transition occurs at the value λ_*^u of the gauge coupling constant λ which becomes smaller as we increase M^2/N until it vanishes for $M^2/N \rightarrow \infty$. By definition the double scaling limit in which the fixed ratio M^2/N is taken to infinity corresponds to the model (18). Indeed the partition function (18) is obtained from the partition function $Z[\gamma, \lambda, M^2]$ given by (19) in the limit in which we take $M^2 \rightarrow \infty$ first and then $N \rightarrow \infty$. Hence we can immediately conclude that the model (18) does not undergo the above one-plaquette phase transition since $\lambda_*^u \rightarrow 0$ when $M^2/N \rightarrow \infty$ and furthermore this model according to the restriction (25) exists mostly in the matrix phase.

Thus the model (24) with finite values of the ratio M^2/N behaves as an ordinary $2d$ lattice gauge model with a reduced one-plaquette critical value whereas in the limit $M^2/N \rightarrow \infty$ the 3rd order one-plaquette phase transition is completely removed from the model. This limit can be thought of as a way of regularizing the one-plaquette transition in the lattice model.

However for $M^2/N \rightarrow 0$ the above computed critical value $\lambda_*^u = 2$ is not the correct critical value at which the third order phase transition should occur. The model (19) with the fixed ratio M^2/N such that $M^2/N \rightarrow 0$ is a $U(2N)$ one-plaquette model which should be described by the partition function (22) and not by (24). By going through the same steps which led to (30) we obtain for the model (22) the critical value $\lambda_{M*} = 1$ or equivalently

$$\lambda_*^l = \frac{1}{1 + \frac{M^2}{N}} = 1 - \frac{M^2}{N} + O\left(\frac{M^4}{N^2}\right). \quad (38)$$

In above we have used the fact that the partition function (22) should really be used only for very small M^2/N . Clearly for $M^2/N \rightarrow 1$ the critical value $\lambda_*^l \rightarrow 0$ and then it becomes negative for $M^2/N > 1$ and hence the above formula should only be valid in the range $M^2/N \in [0, 1[$. In other words the 3rd order one-plaquette phase transition occurs at the values λ_*^l of λ for all $M^2/N \in [0, 1[$ whereas for $M^2/N \in [1, \infty[$ the transition should occur at the values λ_*^u . The value $M^2/N = 1$ seems to demarcate a discontinuity in the phase diagram. Below $M^2/N = 1$ the model (19) describes an ordinary $U(2N)$ gauge theory on one plaquette whereas above this value the model becomes the $U(N)$ gauge theory on two plaquettes given by (24) which corresponds to noncommutative gauge theory on the fuzzy sphere. The numbers of degrees of freedom of the model below and above this critical mass $M^2/N = 1$ are different. Above $M^2/N = 1$ we have $2N^2$ degrees of freedom whereas below this value we have $4N^2$ degrees of freedom.

Thus the model (19) with $M^2/N = 1$ is seen to be the first noncommutative $U(1)$ gauge model on the fuzzy sphere S_N^2 which we will obtain as we increase M^2/N from 0 to ∞ . Indeed this model has the correct number of degrees of freedom given by $2N^2$ and it can be reduced to (24). The model with $M^2/N = 1$ is also the noncommutative model with the largest “fuzzy sphere” phase in the sense of (25) hence the effects of the “matrix” phase are the weakest in this case. As a consequence the estimation of the critical point of the 3rd order phase transition is




Figure 1: The weak-coupling and strong-coupling phases of the noncommutative $U(1)$ gauge theory on the fuzzy sphere are the regions of the weak-coupling and strong-coupling phases with $M^2/N \geq 1$. The regions with $M^2/N < 1$ correspond to an ordinary $U(2N)$ gauge theory. The number of degrees of freedom of the model above the critical mass $M^2/N = 1$ is $2N^2$ whereas below $M^2/N = 1$ we have $4N^2$ degrees of freedom. The fuzzy sphere phase consists of the weak-coupling and strong-coupling phases with $M^2/N \geq 1$. In this phase the theory is a $U(N)$ gauge theory on two plaquettes.

most reliable using the model (19) with $M^2/N = 1$. At $M^2/N = 1$ the critical value λ_*^u is given by $\lambda_*^u = 2/3 = 0.66$. This computed critical value leads to the critical value of the coupling constant $\bar{\alpha}$ (from equation (16))

$$\bar{\alpha}_*^4 = \frac{64}{\lambda_*^u} = 96 \Leftrightarrow \bar{\alpha}_* = 3.13. \quad (39)$$

This is to be compared with the observed value $\bar{\alpha}_* = 3.35 \pm 0.25$ seen in Monte Carlo simulation of the model (1) with the Metropolis algorithm. The agreement is very good. Thus it seems that the two noncommutative $U(1)$ models given by (1) (with large m^2) and (19) (with $M^2/N = 1$) are quantum mechanically equivalent in the fuzzy sphere S_N^2 phase.

4 Conclusion

We constructed in this article a one-parameter family of noncommutative $U(1)$ gauge theory partition functions $Z_M \equiv Z(\gamma, \lambda, M^2)$ on the fuzzy sphere S_N^2 given by equation (19). The idempotent γ is the chirality operator and λ is the gauge coupling constant. In the limit $M^2 \rightarrow 0$ the model becomes a $U(2N)$ one-plaquette model whereas in the limit $M^2 \rightarrow \infty$ we

get the generalized $U(N)$ one-plaquette model defined by (18). In the double scaling limits $M^2, N \rightarrow \infty$ keeping M^2/N fixed ≥ 1 we can show that quantum noncommutative $U(1)$ gauge theory on the fuzzy sphere S_N^2 is equivalent to a $U(N)$ nonabelian Yang-Mills theory on a two dimensional lattice with two plaquettes according to (24). Thus the model (19) (with $M^2/N = 1$) will undergo in the fuzzy sphere phase a 3rd order one-plaquette large N phase transition between weak-coupling and strong-coupling phases around the critical value (39) which is consistent with the observed 3rd order transition point of (1). In the “matrix” phase we will have a $U(N)$ gauge theory on a single point.

Another evidence for this equivalence comes from the calculation of the specific heat which seems to agree with the simulation results of (1) both in the weak-coupling and strong-coupling phases up to the sphere-to-matrix first order transition where the whole spacetime (the sphere) collapses. In particular the specific heat was found to be equal to 1 in the fuzzy sphere weak-coupling phase of the gauge field which agrees with the observed value 1. The value 1 comes precisely because we have two plaquettes which approximate the noncommutative $U(1)$ gauge field on the fuzzy sphere. In the strong-coupling region deviations between (37) and the data coming from the simulation of (1) become significant only near the sphere-to-matrix transition point [5, 6].

References

- [1] Badis Ydri, *Mod.Phys.Lett.A* 19 (2004)2205-2213, *Nucl.Phys.B* 690 (2004)230-248.
- [2] J.Madore, *Class.Quant.Grav.* 9:69-88,1992. J.Hoppe, MIT PhD thesis (1982).
- [3] A.Y.Alekseev , A.Recknagel, V.Schomerus, *hep-th/0003187* and *hep-th/9812193*.
- [4] S.Iso, Y.Kimura, K.Tanaka, K.Wakatsuki, *Nucl.Phys.* B604 (2001) 121.
- [5] D.O’Connor, Badis Ydri , *Monte Carlo Simulation of NC Gauge Field on the Fuzzy Sphere*, *hep-lat/0606013*.
- [6] Badis Ydri , *The one-plaquette model limit of NC gauge theory in 2D*, *hep-th/0606206*.
- [7] P.Castro-Villarreal , R.Delgadillo-Blando , Badis Ydri, *hep-th/0405201* , *Nucl.Phys.B* **704** (2004) 111-153.
- [8] T.Azuma,S.Bal,K.Nagao,J.Nishimura,*JHEP* 0405 (2004) 005.
- [9] H.Steinacker, *Nucl.Phys.B*679,66 (2004). See also the recent preprint : H.Steinacker, R.J.Szabo, *hep-th/0701041*.
- [10] D.J.Gross, E.Witten, *Phys.Rev. D* 21 (1980)446-453. S.R.Wadia, EFI preprint EFI 80/15 (March 1980), *Phys. Lett. B* **93** (1980) 403.